



syngenta®

Reg. No. L8914 Act / Wet No. 36 of / van 1947

SAADBEHANDELINGSGROEP 4 SEED TREATMENT GROUP

Aktiewe bestanddeel / Active ingredient:  
**Mefenoxam (fenielamiede) /**  
**Mefenoxam (phenylamide) ..... 240 g/l**

UN Not classified / Nie geklassifiseer

Product names marked ® or ™, the ALLIANCE FRAME  
 the SYNGENTA Logo and the PURPOSE ICON  
 are Trademarks of a Syngenta Group Company.

'n Vloeibare konsentraat sistemiese swamdoder as 'n deurdrenkingsbehandeling vir verwelksiekte op groente, sierplante en bosbou saailinge en as 'n blaarbespuiting op turf.

A flowable concentrate systemic fungicide as a drench treatment for damping off in vegetable, ornamental and forestry seedlings and as a foliar application on turf.

EMERGENCY TEL. / NOOD TEL.  
 ++27 (0) 8312 33 911  
 (Bateleur 911)

© SYNGENTA, AG

Registration holder / Registrasiehouer:  
**Syngenta SA (Pty) Ltd. / (Edms) Bpk.**  
 (Co./Mpy Reg. No. 1998/013761/07)  
 Private Bag / Privaatsak X 60  
 HALFWAY HOUSE, 1685 RSA.  
 Tel. (011) 541-4000

Batch No

Date of Manufacture



HARMFUL SKADELIK



## 1. WARNINGS

- Harmful if swallowed.
- Toxic to daphnia (scale animals) and animals
- Store in a cool dry place.
- Store away from food and feed.
- Avoid storage above 35 °C.
- Keep out of reach of children, uninformed persons and animals.
- Do not use treated seed for human or animal consumption.
- **In case of poisoning call a doctor and make this label available to him.**

Although this remedy has been extensively tested under a large variety of conditions, the registration holder does not guarantee that it will be effective under all conditions. The activity and effect may be affected by factors such as abnormal soil, climatic and storage conditions, quality of dilution water, compatibility with other substances not indicated on the label and the occurrence of resistance of the pathogen against the remedy, as well as by the method, time and accuracy of application. The registration holder furthermore does not accept responsibility for damage to crops, vegetation, the environment or harm to man or animal, or for lack of performance of the remedy concerned due to failure of the user to follow the label instructions, or to the occurrence of conditions which could not have been foreseen in terms of the registration. Consult the supplier in the event of any uncertainty.

## 2. PRECAUTIONS

- Avoid contact with skin, eyes or clothing.
- Do not inhale concentrate or mixed product.
- Wash contaminated clothing after use.
- Do not eat, drink or smoke whilst mixing or applying the product or before washing hands and face.
- While handling the concentrate, during treatment of the seed and while handling treated seed wear a face shield, hat or cap and rubber gloves.
- Clean applicator after use. Dispose of rinsate where it will not contaminate crops, pastures, rivers, dams and boreholes.
- Prevent contamination of food, feed, drinking water and eating utensils.
- Destroy empty container and do not use it for any other purpose.
- When treated seed is stored in containers, mark containers clearly "POISONED" and never use them for any other purpose.

## FIRST AID TREATMENT

**General:** Place the patient in a well-ventilated area and protect him from hypothermia. IN CASE OF SUSPECTED POISONING: Immediately call a physician.

**Eye contact:** Rinse eyes with a light stream of clean water for 10 - 15 minutes.

**Ingestion:** Repeatedly administer medical charcoal in a large quantity of water.

NOTE: Never give anything by mouth to an unconscious person. Do not induce vomiting.

**Skin contact:** Remove contaminated clothing and thoroughly wash the affected parts of the body with soap and water.

**Inhalation:** Immediately move patient to fresh air.

## NOTE TO PHYSICIAN:

Antidote: No specific antidote is known. Apply symptomatic therapy.

## 3. RESISTANCE MANAGEMENT

**SUBDUE MAXX** is classified as a group code 4 fungicide. Any fungus population may contain individuals naturally resistant to **SUBDUE MAXX** and other group code 4 fungicides. The resistant individuals can eventually dominate the fungus population if these fungicides are used repeatedly and exclusively in programs. These resistant fungi may not be controlled by **SUBDUE MAXX** or any other group code 4 fungicides.

To delay fungicide resistance:

- Avoid exclusive repeated use of fungicides from the same fungicide group code. Alternate or tank mix with products from different fungicide group codes.
- Refer to individual product labels when alternating products or when using tank mixtures with products in fungicide group codes 7, 27, 28, 29, 30 and M.
- Integrate other control methods (chemical, cultural, biological) into disease control programs.
- For specific information on resistance management, contact the registration holder of this product (Syngenta South Africa) or visit the FRAC website at <http://www.frac.info/frac/index.html>.

## 4. PRODUCT PROPERTIES

**SUBDUE MAXX** contains mefenoxam, a systemic fungicide with protective and curative properties.

## 5. DIRECTIONS FOR USE Use only as indicated

### 5.1 COMPATIBILITY

The compatibility of **SUBDUE MAXX** with other products may be influenced by the formulation of the products involved as well as the quality of the dilution water. Since the formulation of other products may change without the knowledge of Syngenta and the quality of water may vary from farm to farm, a physical compatibility test should always be carried out prior to application.

## 5.2 RECOMMENDATIONS

DISEASE	CROP/DOSAGE
<b>Damping-off</b> <i>Pythium</i> spp.	<b>VEGETABLE, ORNAMENTAL AND FORESTRY SEEDLINGS</b>  7.7 ml/15 ℓ water  Apply as a drench to seedling trays using 1.5 ℓ of mixture/m <sup>2</sup>
<b>REMARKS:</b> <ul style="list-style-type: none"> <li>Applications of <b>SUBDUE MAXX</b> should commence when seedlings are at the 3 - 4 leaf stage and repeated every 14 days until transplant.</li> <li>No more than three (3) applications per seedling cycle may be done.</li> </ul>	
<b>Pythium leaf blight, Pythium root rot</b> and <b>Damping-off</b> <i>Pythium</i> spp.	<b>TURF</b>  1.7 to 3.5 ℓ/ha or 17 to 35 mℓ/100 m <sup>2</sup>
<b>REMARKS:</b> <ul style="list-style-type: none"> <li>Apply <b>SUBDUE MAXX</b> as part of a preventative fungicide programme containing fungicides from different chemical groups.</li> <li>Apply before symptoms occur when condition favours disease development.</li> <li>Apply no more than two (2) consecutive <b>SUBDUE MAXX</b> applications at 14 to 21 day intervals.</li> </ul>	

**SUBDUE MAXX**<sup>®</sup> = Registered Trade mark of a Syngenta Group Company.  
**MEFENOXAM**<sup>™</sup> is the trade name of metalaxyl-M (ISO common name).

© SYNGENTA AG, 2000. Copyright in this document is reserved.  
All unauthorized reproduction is forbidden.

SA Reg.: 2013-07-04